

High Risk Meningococcal Vaccine Order Form

Please complete form and fax to (807) 625-4828.

Client: _____ D.O.B.: _____ Health Card #: _____
(yyyy/mm/dd)

Clinic: _____ Health Care Provider: _____

Eligibility for the high risk meningococcal vaccine includes those who have one or more of the Following medical conditions:

- Functional or anatomic asplenia
- Complement, properdin, factor D deficiency or primary antibody deficiencies
- Cochlear implant recipients (pre/post implant)
- Acquired complement deficiency
- HIV

VACCINE REQUESTED:

Menactra (Men-C-ACYW)

A. Persons between 9 months to 55 years of age and who have one or more of the medical conditions listed above are eligible to receive age appropriate primary series with Men-C-ACYW-135 (Menactra) vaccine and booster doses. See table 1 for more details.

Table 1: Men-C-ACYW (Menactra)

Age	Recommended Intervals
9 to 11 months	<ul style="list-style-type: none"> ▪ 1st dose given ▪ 2nd dose, 2 months after 1st dose ▪ 3rd dose, 2 months after 2nd dose ▪ Booster doses every 5 years
12 months to 6 years	<ul style="list-style-type: none"> ▪ 1st dose given ▪ 2nd dose, 2 months after 1st dose ▪ Booster every 3 to 5 years
7 to 55 years	<ul style="list-style-type: none"> ▪ 1st dose given ▪ 2nd dose 2 months after 1st dose ▪ Booster doses every 5 years

Menomune (Men-P-ACYW)

B. Persons older than 55 years and who have one or more of the medical conditions listed above are eligible to receive one dose of Men-P-ACYW (Menomune) vaccine. Menomune should not be used for booster doses.

Bexsero (4CMenB)

C. Persons 2 months through 17 years of age with one or more of the medical conditions listed above are eligible to receive the meningococcal B vaccine or Bexsero. See Table 2 for more details.

Table 2: 4CMenB (Bexsero)

Age at first dose	Doses	Recommended Intervals
2 to 5 months	3+1	<ul style="list-style-type: none"> ▪ 1st dose given ▪ 2nd dose, 2 months after 1st dose ▪ 3rd dose 2 months after 2nd dose ▪ 4th dose booster, 2 months after 2nd dose and at age \geq 12 months
6 to 11 months	3	<ul style="list-style-type: none"> ▪ 1st dose given ▪ 2nd dose, 2 months after 1st dose ▪ 3rd dose, 2 months after 2nd dose and at age \geq 12 months
12 months to 10 years	2	<ul style="list-style-type: none"> ▪ 1st dose given ▪ 2nd dose 2 months after 1st dose
11 to 17 years	2	<ul style="list-style-type: none"> ▪ 1st dose given ▪ 2nd dose, 1 month after 1st dose

Notes:

- The vaccines are available to close contacts of a case if IMD and during outbreaks caused by N. meningitis serogroups A,C,Y,W and B
- For more details, please see the Publicly Funded Immunization schedule for Ontario for recommended schedule and number of doses by age.

IMT-221 (2016) page 1 of 1