

# TICKS

## PREVENT THE BITE

- Cover up as much skin as possible
- Use bug spray with Icaridin or DEET on exposed skin
- Check yourself and your pets before heading inside

## REMOVE THE TICK

- Use fine-tipped tweezers to grab as close to the skin as possible
- Pull up slowly, don't twist or turn
- Wash with soap OR apply rubbing alcohol
- Monitor for unusual symptoms; see a health care provider if concerned

## SUBMIT THE TICK

- Place in a clean container and submit to the Health Unit



Blacklegged tick in various stages of feeding compared to the size of a dime. (Source: Public Health Agency of Canada)

The City of Thunder Bay and immediately surrounding area are now considered a risk area for Lyme disease. About 10% of the blacklegged ticks in this area are infected with Lyme disease bacteria.

These bacteria can be transmitted to humans after a female blacklegged tick has fed for 24 or more hours.

Most ticks found in the Thunder Bay District are wood ticks and do not carry the Lyme disease bacteria.

rev. May 2019

[TBDHU.COM/ticks](http://TBDHU.COM/ticks) for more info

# TICK IDENTIFICATION CARD

## Blacklegged (Deer) Ticks



**Adult Male**

All black



**Adult Female**

Reddish-brown body/black cape and legs

## Wood (Dog) Ticks



**Adult Male**

Mottled brown-cream back/reddish-brown legs



**Adult Female**

Brown body/cream cape/reddish-brown legs

Photo credit: University of Rhode Island TickEncounter Resource Centre