

Ministry of Health

Chapter 1: Storage and Handling of Pfizer-BioNTech's COVID-19 Vaccines

Version 8.0 – October 22, 2024

Highlights of Changes:

- Update to current KP.2 formulation

The scope for this chapter includes information pertaining to the storage and handling of Pfizer-BioNTech's COVID-19 Vaccine.

The intended audience for this guidance document includes all providers that are:


- Storing, distributing and/or administering COVID-19 vaccines;
- Involved in the assessment of temperature excursions, including the vaccine return process;

Additional resources available:

- Pfizer-BioNTech: <https://www.cvdvaccine.ca/>
- [COVID-19 Vaccine Storage and Handling Guidelines](#)
- [Vaccine Storage Handling Guidelines](#);
- Individual product monographs on the [Government of Canada website](#).

In addition, health care providers and organizations who have questions should contact their [local public health unit](#) or the Ministry of Health at vaccinesupplyandlogistics@ontario.ca.

Table 1. Pfizer-BioNTech formulations of the COMIRNATY® vaccine available in Ontario

Grey Cap
KP.2 Formulation

30 mcg/0.3mL 6 doses (do not dilute)

Storage and Handling of Pfizer-BioNTech's COVID-19 Vaccines

Table 2. Storage and Handling for Pfizer-BioNTech (KP.2 Formulation)

Storage Condition	Grey Cap
Frozen Vials Prior to Use	<ul style="list-style-type: none"> • Store at ultra-low temperature freezer at -90°C to -60°C until printed expiry date or otherwise noted. • Do not store vials at -25°C to -15°C. • Keep protected from light, in the original packaging, until ready to use.
Thawed Unpunctured Vials	<ul style="list-style-type: none"> • Thawed vials may be stored in the fridge between +2°C to +8°C for a single period of up to 10 weeks. The 10-week refrigerated expiry date should be recorded on the carton at the time of transfer. • Vaccine may be stored at room temperature +8°C to +25°C up to a maximum of 12 hours. • Thawed vials can be handled in room light conditions. • Do not refreeze thawed vials.
Thawed Punctured Vials or Syringes	<ul style="list-style-type: none"> • After first puncture, the vial should be stored between +2°C to +25°C. • Vials should be discarded 12 hours post-puncture. • If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and any excess volume. Do not pool excess vaccine from multiple vials.

Rounding Principles

Based on information from Pfizer-BioNTech; COVID-19 vaccines at refrigerated temperatures may be rounded to the nearest whole degree:

- Temperatures between +1.5°C and +1.9°C are rounded to +2.0°C
- Temperatures between +8.1°C and +8.4°C are rounded to +8.0°C

Pfizer BioNTech COVID-19 vaccines exposed to temperatures between +1.5°C and +8.4°C are considered to be in refrigerated temperatures and the incident does not need to be recorded as a temperature excursion and entered in COVAX_{ON}. Troubleshooting should occur to ensure that temperatures are corrected and maintained between +2°C to +8°C.

Thawing Pfizer-BioNTech Vaccine

Table 3. Thawing Conditions for Pfizer BioNTech’s COVID-19 Vaccine

Thawing Condition	Grey Cap
Thawing in a refrigerator	<ul style="list-style-type: none"> • Thaw in the refrigerator at +2°C to +8°C. • A carton of 10 vials may take up to 6 hours to thaw and thawed vials can be stored in the refrigerator for up to 10 weeks.
Thawing at room temperature	<ul style="list-style-type: none"> • If not previously thawed at +2°C to +8°C, allow vial(s) to thaw at room temperature (up to +25°C) for 30 minutes

Transport Conditions for Pfizer-BioNTech Vaccines

Table 4. Transport Conditions for Pfizer-BioNTech COVID-19 Vaccines

Storage Condition	Grey Cap
Vaccine During Transport (by vehicle on ground, air, or water)	<ul style="list-style-type: none"> • Transport in ULT. • If ULT not possible, transport between +2°C to +8°C. <ul style="list-style-type: none"> ○ This time counts toward the 10-week storage limit • Do not pack vaccine that is between +2°C and +8°C with frozen vaccine. • Store upright and protect from light. • Label as fragile. • Protect from shocks, drops, vibration, etc. • No data is available regarding the stability of punctured vials or pre- drawn syringes during transportation.

Transportation Of Pfizer-BioNTech COVID-19 Vaccine in Syringes

Whenever possible, it is recommended that Pfizer-BioNTech COVID-19 vaccine be transported in an unpunctured vial and that the entire vial be administered in one location rather than transporting syringes filled with vaccine.

However, while not suggested as routine practice, in exceptional circumstances, diluted (if applicable) vaccine maybe transported in a syringe.

- The vaccine does not contain a preservative, therefore special attention should be given to handling and packaging of the syringe to prevent contamination.

Exceptional circumstances may include situations in which a few doses are needed to support the immunization and series completion of small numbers of individuals residing in congregate settings (i.e., one or two residents) and for those who are home bound (e.g., those who may be unable to attend a community-based clinic due to physical limitations).

Example of pre-drawn syringe and container labels:

Pfizer-BioNTech COMIRNATY COVID-19 Vaccine <insert age indication>

Facility name and phone number:

Quantity of syringes:

Date prepared & Time to discard:

Lot #:

Initials of preparer:

Vaccine Transportation Scenarios

The following scenarios may assist with planning the transportation of vaccines.

Scenario 1: Ground Transport between Locations

Transporting vaccine from a hospital to another hospital for longer term storage (ULT or frozen).

Transportation of the vaccine for storage at another facility should be done in the frozen state at ultra-cold temperature.

Scenario 2: Ground Transport between Locations or Facilities

Transporting vaccine from a hospital to a congregate living setting.

Transportation of the vaccine should be carried out in ULT, frozen state or liquid state transport. See product monograph and above details and recommendations.

Scenario 3: Medium and Long Duration Ground PLUS Air Transport

At this time, it is recommended that any transport that involves air be done in a frozen state, ultra-cold temperature, per product monograph details and recommendations above.

Scenario 4: Short Duration Movement within a Facility or Campus

Movement of the vaccine that is stored at a long-term care home but needs to be walked over to an attached retirement home (e.g., on the same campus/property) can be transported using a Playmate cooler and well-functioning wheeled cart on a relatively smooth pathway. Transport may also be conducted as a hand-carry (walked only, no running). Following general precautions described above, such movement may be conducted for a short period.

Vaccine storage post temperature excursion or unit malfunction.

Vaccines placed in an ULT portable unit (-90°C to -60°C) can go back into an ULT unit. To the extent possible, vials should be kept in the tray during transport. If this is not possible, the vials need to be securely stored (not rolling around) in the storage device.

- If vaccines are placed in an insulated container for +2°C to +8°C temperature range, the vaccines should go back into a refrigerator and not be refrozen.
 - Time in fridge temperatures counts toward the 10-week storage limit

If the purpose-built vaccine storage unit is unable to maintain the required storage temperature range, maintain the vaccines in the assigned container and continue to monitor temperatures inside the container. Place the vaccine back into the purpose-built unit once it is able to maintain the temperature range as specified by the vaccine manufacturer in the product monograph.

For manufacturer contact information please see the General COVID-19 Vaccine Storage and Handling Guidance on the [COVID-19 Vaccine Program](#) website.