



West Nile Virus in the Thunder Bay District, 2025




Dr. Ken Deacon
Deacon Bioconsulting



Prepared for the Thunder Bay District Health Unit



2026



Thunder Bay District Health Unit

West Nile Virus in the Thunder Bay District, 2025

Table of Contents

List of Figures	3
List of Tables	3
Executive Summary	4
History of West Nile Virus in North America	6
Illness Caused by West Nile Virus	7
Transmission of West Nile Virus in Humans	7
Mosquito Vectors	7
Objectives of the WNV Surveillance Program, 2025	9
Larval Mosquito Surveillance	10
Adult Mosquito Surveillance	10
Bird Surveillance	15
Mammal Surveillance	15
Human Surveillance	15
West Nile Virus Control Measures	15
Conclusions	16
Recommendations for 2026	17
Literature Cited	18

Visit us at www.tbdhu.com

List of Figures

- Fig. 1 Location of CDC adult mosquito light traps in the city of Thunder Bay and surrounding area, 2025. 11
- Fig. 2 Number of individuals of male, unidentifiable, non-vector and vector mosquito species collected in the city of Thunder Bay and surrounding area, 2025. 12
- Fig. 3 Number of individuals of vector mosquito species collected in the city of Thunder Bay and surrounding area, 2025. 13

List of Tables

- Table 1a Human infections and deaths in Canada from West Nile virus, 2002-2012. 6
- Table 1b Human infections and deaths in Canada from West Nile virus, 2013-2025. 6

EXECUTIVE SUMMARY

No human case of West Nile Virus (WNV) was reported from the District of Thunder Bay during 2025. Three birds tested positive for West Nile virus. Mosquitoes collected in Centers for Disease Control and Prevention (CDC) adult mosquito light traps provided the information necessary to evaluate human risk from WNV in 2025.

Twelve CDC adult mosquito light traps were operated for one night per week in the city of Thunder Bay and the surrounding area on 24 June and during the eight-week period from 8 July to 26 August. Unfortunately, dry ice, which is crucial for the operation of the traps, was unavailable on two occasions, namely 24 June (week 26) and 19 August (week 34). The CDC traps were operated on these two occasions using only the ultra violet lights and fans; therefore, fewer mosquitoes were collected than with the usual dry ice. Mosquitoes were collected with dry ice on 84 trap-nights and without dry ice on 24 trap nights. The partial loss of data from week 26 and week 34 marginally compromised the risk analysis for 2025. Twelve vector species (three enzootic and nine epizootic) were found in the light traps during the summer of 2025. The light traps collected a total of 10348 individuals. A total of 4302 individuals were identified (not including males and damaged specimens) from the 10348 individuals collected in the light traps. The identified individuals contained 1835 vectors. The vectors were pooled by species, then 43 pools of *Culex restuans*, 11 pools of *Ochlerotatus japonicus* and one pool each of *Culex salinarius*, *Culex tarsalis*, *Aedes vexans* and *Ochlerotatus stimulans* were tested for WNV using Reverse Transcription-Polymerase Chain Reaction (RT-PCR). No mosquito pool tested positive for WNV. *Culiseta melanura*, the principal vector of Eastern Equine Encephalitis Virus (EEEV) was not collected during 2025; therefore, no tests for EEEV were completed.

An estimate of the total number of each species of the vector mosquitoes was calculated (n= 2146). The three enzootic vectors found were *Culex restuans* (n= 268), approximately 2.59%, *Culex salinarius* (n= 1) and *Culex tarsalis* (n= 1) comprised less than 0.01% of the 10348

mosquitoes. The nine epizootic vectors included *Aedes vexans* (n= 1550) (14.98%), *Ochlerotatus canadensis* (n= 103) (1.00%), *Ochlerotatus triseriatus* (n= 94) (0.91%) *Oc. stimulans* (n= 54) (0.52%), and *Oc. japonicus* (n= 23) (0.22%). Five other epizootic species comprised less than one percent each of the mosquitoes collected.

A value of only 67.3 accumulated degree-days (AccDD) was recorded in Thunder Bay from 1 June to 31 August during 2025. A value of 380 AccDDs is required before there is a serious risk of a WNV outbreak.

The low incidence of the enzootic vector *Cx. restuans* (2.59%), the probable absence of *Culex pipiens*, as well as the low AccDD value made the amplification of WNV within the bird community to the point of “spill-over” an extremely unlikely event during the summer of 2025. The risk to humans of acquiring WNV was considered minimal. Neither larviciding nor adulticiding was considered necessary during 2025.

If the climate of Thunder Bay changes to one which is more similar to the prairies (hot and dry) or to that of southern Ontario (hot and humid), there will be increased risk from WNV because of increases of *Cx. restuans*, *Culex tarsalis*, *Cx. salinarius*, *Oc. japonicus* and perhaps the establishment of *Culex pipiens*. Any increase in these extremely competent vectors would increase the risk of an outbreak of WNV in the District of Thunder Bay.

History of West Nile Virus in North America

West Nile virus (WNV), a mosquito-vectored disease, first appeared in North America in New York City during 1999 (Marr & Calisher 2003). Subsequently WNV spread across the continent. As of 13 January 2026, all of the continental states except for Alaska, Vermont and New Hampshire reported human WNV infections (Centers for Disease Control and Prevention 2026). A total of 2076 cases of WNV infections occurred in humans in the United States during 2025 (Centers for Disease Control and Prevention 2026).

WNV was first detected in Canada during 2001 when dead birds tested positive for the virus in southern Ontario (Public Health Ontario 2016). The first human cases of WNV in Canada occurred during 2002 in Ontario and Quebec (Public Health Ontario 2016). One peak in the incidence of infections (n=1481) occurred during 2003 (Table 1a) when the disease spread across eastern Canada. A second peak (n=2215) occurred during 2007 (Table 1a) when the disease entered the prairie region of Canada. WNV Cases reported up to 1 November 2025 were 295 of which 114 were from Quebec, 169 from Ontario, 3 from Manitoba, 3 from Saskatchewan, 5 from Alberta and 1 from British Columbia (Table 1b). Nine deaths were attributed to WNV in Canada during 2025 (Table 1b).

Table 1a. Human infections and deaths in Canada from West Nile virus, 2002-2012 (Government of Canada 2025).

	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Clinical Cases	414	1481	25	225	151	2215	36	13	5	101	428	115
Deaths	11	10	2	12	2	8	0	0	0	4	6	4

Table 1b. Human infections and deaths in Canada from West Nile virus, 2013-2024 (Government of Canada 2025).

	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025
Clinical Cases	21	80	104	200	367	37	86	35	27	75	167	295
Deaths	0	0	7	8	26	0	3	0	0	2	0	9

Illness Caused by West Nile Virus

WNV is vectored by infected mosquitoes. Eighty percent (80%) of people infected with WNV are asymptomatic. Twenty percent (20%) of infected people develop West Nile fever which consists of fever, tiredness, headache, muscle aches, rash and/or swollen glands.

Approximately one in 150 people infected with the virus will develop a life-threatening manifestation called encephalitis, *i.e.* a swelling of the brain. Symptoms of West Nile encephalitis include fever, headache, stiff neck, disorientation, tremors, muscle weakness, paralysis and/or coma (Elliott *et al.* 2003). West Nile encephalitis is more common, but not restricted to people over 50 years of age. The severity of the disease increases with age (Drebot & Artsob 2006). From 4% to 14% of people with West Nile encephalitis will die as a result of their infection, whereas others may experience long-lasting, debilitating problems ranging from memory loss to muscle weakness (Elliott *et al.* 2003).

Transmission of West Nile Virus to Humans

West Nile virus is a mosquito-borne flavivirus which primarily infects birds, producing a transient high viraemia that ensures transmission of the virus back to feeding mosquitoes in an amplifying cycle. The amplification cycle does not occur in mammals. The virus replicates in only a few species of mosquitoes which act as the vectors of this disease. Humans can become infected as a result of bites from mosquitoes that have bitten infected birds. Other less common routes of transmission include: intrauterine, breast milk, blood transfusions, organ transplants, as well as needle stick or sharps injuries. Immuno-compromised patients and the elderly are at the greatest risk for encephalitis and death (Groner 2005).

Mosquito Vectors

Mosquitoes either overwinter with the virus or become infected with WNV when they bite infected birds. According to Reisen *et al.* (2006), replication of the virus occurs in the mosquito at a temperature above 14.3°C; however, work by Cuevas (in Hunter & Gasparotto 2015)

indicates that the threshold temperature for the development of the virus is 16.3°C. The Ontario Ministry of Health and Long-Term Care uses a value of 18.3°C as the threshold temperature for the calculation of viral replication. Infected enzootic vector mosquitoes bite birds, thus transmitting the infection to new hosts (Elliott *et al.* 2003). The virus undergoes replication in the newly infected birds. These birds then become a source of infection for other mosquitoes as this enzootic phase of the disease progresses in an amplifying cycle. If amplification begins during the early spring, then by mid-summer a large number of infected birds and mosquitoes are present which creates a high risk for human infection.

The primary mosquitoes involved in the enzootic amplification process in southern Ontario are *Culex pipiens* and *Culex restuans*. These mosquitoes prefer to feed on birds but may also bite humans or other mammals (Wood *et al.* 1979). *Cx. pipiens* is now known to be attracted to humans at certain times during its lifecycle which means that this species also acts as an epizootic vector of WNV to humans (Russell 2008). Epizootic vectors, also called “bridge vectors”, transmit WNV from birds to mammals. A third *Culex* species, *Culex tarsalis*, is the main mosquito species responsible for the transmission of WNV in western North America (Goddard *et al.* 2002). *Cx. tarsalis* has occasionally been found in the Thunder Bay District (Deacon 2013), but is primarily found in the prairies (Wood *et al.* 1979). *Cx. tarsalis* is a highly competent vector within the genus *Culex* because it feeds freely on both birds and mammals (Wood *et al.* 1979) thus acting as both the enzootic and epizootic vector of WNV. The ability of *Cx. tarsalis* to feed on birds and mammals (especially humans) probably accounts for the magnitude of the WNV outbreak in the prairies during 2007 (Table 1a).

Cx. pipiens and *Cx. restuans* prefer to lay their eggs in man-made structures that contain water such as street-side catch basins, road-side ditches, and man-made containers (tires, bottles, buckets, bird baths, roof gutters, swimming pool covers, rain barrels, etc.) where the eggs develop into larvae, then pupae and finally adults (Wood *et al.* 1979). *Cx. tarsalis* lays its eggs in permanent and semi-permanent ponds, irrigation and roadside ditches with emergent vegetation, as well as temporary pools or artificial containers (Wood *et al.* 1979).

“Bridge vectors” (epizootic vectors) are generalist feeders, biting both birds and mammals. “Bridge vectors” are responsible for transmitting WNV from birds to humans during a “spill-over” (the epizootic phase of the disease) which occurs during the late summer. The “spill-over” occurs only when a large number of infected birds are present. Currently *Aedes vexans* is the principal “bridge vector” in Ontario; however, this species is only moderately effective as a vector of WNV (Turell *et al.* 2001). *Ae. vexans* breeds in temporary pools, marshes, and swamps (Wood *et al.* 1979) and is abundant in the District of Thunder Bay. *Ochlerotatus canadensis* is another “bridge vector” species which is sometimes abundant in the District of Thunder Bay. Larvae are found in small open ponds, temporary woodland pools, roadside ditches, cattail and sedge marshes, and muskeg pools (Wood *et al.* 1979). *Oc. canadensis* is considered a moderately effective “bridge vector” of WNV (Belton 2007).

Cx. pipiens, *Cx. restuans*, *Cx. tarsalis*, *Ae. vexans*, and *Oc. canadensis* are found in close proximity to human populations which makes these mosquitoes important in the transmission of WNV. Both enzootic and epizootic vectors are required in high numbers near humans before WNV can be transmitted to humans.

Objectives of the West Nile Virus Surveillance Program, 2025

1. A risk analysis of West Nile Virus activity in the Thunder Bay District was to be completed.
2. Adult mosquitoes in the District of Thunder Bay were to be collected using Centers for Disease Control (CDC) adult mosquito light traps.
3. Adult mosquitoes were to be identified to species and the prevalence of West Nile Virus in vector species was to be determined using Reverse Transcription-Polymerase Chain Reaction (RT-PCR).
4. The habitat used by *Cx. tarsalis* in the District of Thunder Bay was to be identified, if possible.
5. Larval mosquito habitat was to be identified and inspected in the city of Thunder Bay.
6. Human cases of WNV within the District of Thunder Bay were to be reported.

7. Geographic Information Systems (GIS) mapping was to note:
 - Mosquito species distributions
 - Larval habitat locations
 - Catch basin data
 - High-risk locations
8. All municipalities within the District of Thunder Bay were to be offered a West Nile Virus information presentation outlining the TBDHU Action Plan and research findings, if desired.
9. Science-based information was to be used to determine the need for chemical control of larval and/or adult mosquitoes.
10. The 2025 report on West Nile Virus activity in the District of Thunder Bay was to be completed.

Larval Mosquito Surveillance

No complaint about potential larval mosquito habitat in the form of standing water was received during 2025. Larval mosquito habitat sampling should continue as necessary in 2026.

Adult Mosquito Surveillance

Twelve CDC adult mosquito light traps were to be operated for one night per week for nine weeks in the city of Thunder Bay and the surrounding area on 24 June and during the eight-week period from 8 July to 26 August. Surveillance did not occur on 1 July (Canada Day), because the live specimens could not be shipped safely for identification and viral testing by Entomogen Inc. Unfortunately, dry ice, which is crucial for the operation of the traps, was unavailable on two occasions, 24 June (week 26) and 19 August (week 34). The CDC traps were operated on these two occasions using only the ultra violet lights and fans; therefore, fewer mosquitoes were collected than with the usual dry ice. Mosquitoes were collected with dry ice on 84 trap-nights and without dry ice on 24 trap nights. The partial loss of data from week 26 and week 34 marginally compromised the risk analysis for 2025. Eleven of the twelve traps were in the same fixed, secure locations as 2024 (Fig. 1). One trap was moved to a new location.

The contents of the light traps were analysed by Entomogen Inc. All the mosquitoes in each trap were identified to species unless the trap contained more than 150 individuals. These larger samples were counted and sub-sampled with at least 150 individuals identified randomly.




Fig. 1 Location of CDC adult mosquito light traps in the city of Thunder Bay and surrounding area, 2025.

The remaining mosquitoes were referred to as “extras”. The light traps collected “extras” on ten occasions during the 2025 trapping season. Entomogen Inc. also performed the viral analyses of selected vector mosquitoes as directed by Public Health Ontario (2014).

Twelve vector species (three enzootic and nine epizootic) were found in the light traps. The light traps collected a total of 10348 specimens. A total of 4302 individuals were identified from these specimens, of which 1835 were vectors. An additional 28 individuals were damaged or unidentified males which do not feed on blood (they do not transmit WNV). All other specimens of the 4302 mosquitoes were identified. The vectors were pooled by species, then 43 pools of *Culex restuans*, 11 pools of *Ochlerotatus japonicus* and one pool each of *Culex salinarius*, *Culex tarsalis*, *Aedes vexans* and *Ochlerotatus stimulans* were tested for WNV using Reverse Transcription-Polymerase Chain Reaction (RT-PCR). No mosquito pool tested positive

for WNV. *Culiseta melanura*, the principal vector of Eastern Equine Encephalitis Virus (EEEV) was not collected during 2025; therefore, no tests for EEEV were completed.

An estimated number of each species of the unidentified specimens was based on the ratio of the individuals identified to the total number of individuals in the light traps per sampling event. The number of identified individuals of each species was multiplied by this ratio to obtain an estimate of the approximate total number of individuals of each species collected on that night. After this conversion, males and damaged individuals numbered 30, non-vector species numbered 8172 individuals and vector species numbered 2146 individuals (Fig. 2). These values are a more accurate reflection of the number of each species within the mosquito community throughout the entire summer.

Cx. restuans, an enzootic vector recovered from the light traps, totaled 268 individuals (Fig. 3), about 2.59% of all the mosquitoes collected. The other enzootic vectors were *Culex salinarius* (n= 1) and *Culex tarsalis* (n= 1) each less than 0.01% of the 10348 mosquitoes. The epizootic vectors were *Ae. vexans* (n= 1550) (14.98%), *Anopheles punctipennis* (n= 17) (0.16%), *Anopheles quadrimaculatus* (n= 3) (0.03%), *An. walkeri* (n= 8) (0.07%), *Oc. canadensis* (n= 103) (0.99%), *Oc. japonicus* (n= 23) (0.22%), *Ochlerotatus stimulans* (n= 54) (0.52%) and *Ochlerotatus triseriatus* (n= 94) (0.92%) and *Ochlerotatus trivitattus* (n= 24) (0.23%) (Fig.2). The majority of the non-vector mosquitoes consisted of *Cq. perturbans*, the cattail mosquito (n=7556) (73.02 %) which is an aggressive biter, active at dusk and abundant from July to early August. *Cq. perturbans* does not vector WNV.

Cx. pipiens, one of the main enzootic vectors of WNV in southern Ontario, could possibly be present in the District of Thunder Bay, but has not been positively identified after 20 years of monitoring. Female *Cx. pipiens* and *Cx. restuans* are considered difficult to separate as species; however, the identification of males is possible. Hundreds of *Culex* larvae were reared to the adult stage during each year of this survey. All adult males have been positively identified as *Cx. restuans* only.




Fig. 2 Number of male, unidentifiable, non-vector and vector mosquitoes collected in the city of Thunder Bay and surrounding area, 2025.




Fig. 3 Number of individuals of vector mosquito species collected in the City of Thunder Bay and surrounding area, 2025.

Cx. salinarius is a competent vector of WNV (Andreadis *et al.* 2004) as is *Cx. tarsalis*. Both species were found in the light traps during 2025. These species are widespread in the District of Thunder Bay, but they occur only in low numbers (Deacon 2014). *Oc. japonicus* is another competent vector of WNV (Moberly *et al.* 2005, Andreadis *et al.* 2004, Goddard *et al.* 2002).

Oc. japonicus was found as a low component in the mosquito community from 2017 to 2021 at 0.005%, but increased to over 0.20% from 2022 to 2025 (Deacon 2024), indicating a slightly increased risk of transmission of WNV to humans. Monitoring must continue to follow the changes in the abundance and distribution of *Cx. tarsalis*, *Cx. salinarius* and *Oc. japonicus* to evaluate the potential threat that these species pose to public health.

The Ontario Ministry of Health and Long-Term Care (OMHLTC) now uses accumulated degree-days (AccDD) (based on average daily temperatures above a threshold temperature of 18.3°C) to evaluate the risk of a WNV outbreak. The replication of WNV in mosquitoes depends on ambient temperature. The risk of a serious WNV outbreak occurs only if mosquitoes and the virus experience at least 380 AccDD, although human cases do occur at 100 to 125 AccDD (Public Health Ontario 2013). Based on the OMHLTC threshold temperature of 18.3°C, a value of 67.3 AccDD was calculated from daily maximum/minimum temperatures recorded from 1 June to 31 August 2025 at the Thunder Bay CS station (Environment Canada 2023)). The value of 67.3 AccDD is well below the required 380 AccDD for a serious WNV outbreak; therefore, human cases were considered extremely unlikely.

The low incidence of the enzootic vector *Cx. restuans* (2.59%) and the absence of *Cx. pipiens*, as well as the low AccDD value of 67.3 made the amplification of WNV within the bird community to the point of “spill-over” an extremely unlikely event during the summer of 2025. The risk to humans of acquiring WNV was considered minimal. If the climate of Thunder Bay changes to one which is similar to the prairies (hot and dry) or to that of southern Ontario (hot and humid), there will be increased risk from WNV because of increases of *Cx. restuans*, *Cx. tarsalis*, *Cx. salinarius*, *Oc. japonicus* and perhaps the establishment of *Cx. pipiens*. Any increase in these extremely competent vectors would result in increased risk of an outbreak of WNV in the District of Thunder Bay.

The loss of data from week 26 and week 34 would not have affected the risk analysis for a WNV outbreak stated above. The collection during week 26 (24 June) would have increased the abundance of the vector mosquitoes in the total for the summer. Mosquito numbers drop

precipitously by early August in Thunder Bay. The loss of data from Week 34 (19 August) would have had an almost negligible number of vector mosquitoes because of the continued natural decrease in mosquito numbers by the end of the summer.

Bird Surveillance

Three birds tested positive for West Nile virus which indicates that West Nile virus is still present in Thunder Bay (Entomogen 2025).

Mammal Surveillance

No mammals were submitted for testing during 2025.

Human Surveillance

No human cases of WNV were noted in the District of Thunder Bay during 2025, although two cases were reported during 2007 (Deacon 2008). In 2005 mosquitoes tested positive for WNV to the east of the TBDHU in the Algoma Health District (Entomogen 2005) and one human case of WNV was reported to the west of the TBDHU in the Northwestern Health Unit during 2006 (Northwestern Health Unit 2006). Although the risk of an individual acquiring WNV in the District of Thunder Bay remains low, continued monitoring of the adult mosquito community is necessary and the identification of larval habitat is essential in the event that control measures are necessary in the future.

West Nile Virus Control Measures

Neither larviciding nor adulticiding was required within the Thunder Bay District. WNV control measures in the Thunder Bay District focused on reducing mosquito-breeding sites on private

and municipal property, and on providing information to the public about the prevention of mosquito bites. During 2025 this information was disseminated to the public through the media.

Conclusions

A low potential for an outbreak of WNV in the District of Thunder Bay existed during 2025. *Cx. pipiens*, an extremely competent vector of WNV was not collected. The abundance of *Cx. restuans* in the District of Thunder Bay was so low that WNV posed a minimal risk to human health during 2025; however, *Cx. tarsalis* and *Cx. Salinarius*, as well as *Oc. japonicus* were detected at low numbers. The risk of an outbreak of WNV will increase only when the climate in the District of Thunder Bay changes to one more typical of southern Ontario or western Canada. Any climate change which increases the numbers of the extremely competent vectors *Cx. restuans*, *Cx. tarsalis*, *Cx. salinarius*, *Oc. japonicus* and allow the establishment of *Cx. pipiens* would increase the risk of an outbreak of WNV in the District of Thunder Bay.

More information is required about larval habitat, the adult mosquito community and catch basins if the numbers of enzootic vectors increase in the light traps. Public outreach encouraging the reduction of artificial breeding sites, and personal protection measures to reduce exposure to mosquitoes should continue. A larviciding programme to augment these proposed actions should be considered only if there is a significantly increased risk of human infection by WNV. Contingency plans for pesticide treatment should be considered to prepare for the possibility that changing weather related to global climate change creates conditions more conducive for the transmission of WNV in Thunder Bay.

Recommendations for 2026

1. Complete a risk analysis of West Nile Virus activity in the Thunder Bay District.
2. Continue the adult mosquito surveillance programme within the city of Thunder Bay and surrounding area using CDC light traps.
3. Identify adult mosquitoes to species and determine the prevalence of West Nile Virus in vector species using Reverse Transcription-Polymerase Chain Reaction (RT-PCR).
4. Continue to inspect larval mosquito habitat that has been reported by, and is of concern to the public.
5. Monitor catch basins if the number of enzootic vectors increases in the light traps but only when weather permits.
6. Investigate the biology of *Cx. tarsalis* and *Oc. japonicus*, if weather permits, to determine where these critical species occur in the boreal forest.
7. Report human cases of WNV within the District of Thunder Bay.
8. Use Geographic Information Services (GIS) to note:
 - Mosquito species distributions
 - Larval habitat locations
 - Catch basin data
 - High-risk locations
9. Offer all municipalities within the District of Thunder Bay a West Nile Virus information presentation outlining the TBDHU Action Plan and research findings, if desired.
10. Determine the need for chemical control of larval and/or adult mosquitoes using science-based information.
11. Complete the 2026 report on West Nile Virus activity in the District of Thunder Bay.

Literature Cited

Andreadis, T.G., J.F. Anderson, C.R. Rossbrinck, and A.J. Main. 2004. Epidemiology of West Nile Virus in Connecticut: A five-year analysis of mosquito data 1999-2003. *Vector Borne and Zoonotic Diseases* 4: 360-378.

Belton, P. 2007. *British Columbia mosquitoes as vectors of West Nile virus*.
<http://www.sfu.ca/~belton/summary.pdf>

Centers for Disease Control and Prevention. 2026.
West Nile virus human and non-human activity by county of residence, 2025.
<https://www.cdc.gov/west-nile-virus/data-maps/current-year-data.html>
Last updated 13 January 2026.

Deacon, K. 2024. *West Nile Virus in the Thunder Bay District, 2023*. Thunder Bay District Health Unit, Thunder Bay.
<https://www.tbdhu.com/resource/west-nile-virus-report-2023>

Deacon, K. 2022. *West Nile Virus in the Thunder Bay District, 2021*. Thunder Bay District Health Unit, Thunder Bay.
https://www.tbdhu.com/sites/default/files/files/basic/2023-04/WNV%202021%20Finale_0.pdf

Deacon, K. 2014. *West Nile Virus in the Thunder Bay District, 2013*. Thunder Bay District Health Unit, Thunder Bay.
<http://www.tbdhu.com/NR/rdonlyres/56CE8000-2A85-4BFD-AE64-35B9C202FF0E/0/2013WNVfinalreport.pdf>

Deacon, K. 2013. *West Nile Virus in the Thunder Bay District, 2012*. Thunder Bay District Health Unit, Thunder Bay.
<http://www.tbdhu.com/NR/rdonlyres/E480FC7B-9A8A-4752-AD36-E78CA2B2862F/0/WNV2012finalreport.pdf>

Deacon, K. 2008. *West Nile Virus in the Thunder Bay District, 2007*. Thunder Bay District Health Unit, Thunder Bay.
<http://www.tbdhu.com/NR/rdonlyres/90F2506E-C33B-4287-A32F-496E44839290/0/2008FinalReportDrDeacon.pdf>

Drebot, M.A. and H. Artsob. 2006. West Nile Virus: A pathogen of concern to older adults. *Geriatrics and Aging* 9: 465-471.

Elliott, S.J., M. Loeb, J. Eyles, and D. Harrington. 2003. *Results of a West Nile Seroprevalence Survey, South Oakville, Ontario 2003*. McMaster Institute of Environment and Health, Hamilton.

http://www.mcmaster.ca/mieh/media/WNV_final_report_2003.pdf

Entomogen. 2005. *Mosquito Surveillance: West Nile Virus Surveillance Report Algoma Health Unit 2005*. <http://www.ahu.on.ca/content/healthinfo/index.asp?CategoryID=1545>

Entomogen. 2025. *West Nile Virus Year End Report Thunder Bay*.

Environment Canada. 2025. *Daily Data Report for Thunder Bay CS June 2025*.

https://climate.weather.gc.ca/climate_data/daily_data_e.html?StationID=30682&timeframe=2&StartYear=1840&EndYear=2017&Day=14&Year=2024&Month=6#

Goddard, L. B., A.E. Roth, W.K. Reisen, and T. W. Scott. 2002. Vector competence of California mosquitoes for West Nile virus. *Emerg Infect Dis* 8.

http://wwwnc.cdc.gov/eid/article/8/12/02-0536_article.htm

Government of Canada. 2025. Vector-borne disease surveillance in Canada: Mosquito-borne disease surveillance: Seasonal update.

<https://health-infobase.canada.ca/zoonoses/mosquito/>

Last updated 10 November 2025.

Groner, G. 2005. West Nile Virus: Researchers make inroads into diagnosis and treatments, *Applied Neurobiology* April 2005.

<http://appneurology.com/showArticle.jhtml?articleId=163100345>

Hunter, F.F. and Gasparotto, A. 2015. *West Nile Virus Mosquito Surveillance Report and Notes on Eastern Equine Encephalitis Virus Mosquito Testing, 2015 Thunder Bay District Health Unit*.

Entomogen.

Marr JS, Calisher CH. 2003. Alexander the Great and West Nile Virus Encephalitis. *Emerg Infect Dis*. 9:1599-1603.

<https://doi.org/10.3201/eid0912.030288>

Moberly, S.P., C. Labor, M. McDonough, B. Foster, A. Estes and D.J. Bentfield. 2005. Discovery of an exotic Asian mosquito: *Ochlerotatus japonicus*, (Diptera: Culicidae) in southern Indiana. *Proceedings of the Indiana Academy of Science* 114: 62-64.

Northwestern Health Unit. 2006. *Media Release: Probable Human Case of West Nile Virus*.

<http://www.nwhu.on.ca/alerts/infectious-disease-advisories.php>

Public Health Ontario. 2016. *Vector-Borne Diseases 2015 Summary Report*.
http://www.publichealthontario.ca/en/eRepository/Vector_borne_diseases_Summary_report_2015.pdf

Public Health Ontario. 2014. *Eastern Equine Encephalitis: History and Enhanced Surveillance in Ontario*. Toronto, ON: Queen's Printer for Ontario. ISBN 978-1-4606-4352-5 [PDF]
<https://www.publichealthontario.ca/-/media/documents/E/2014/eeev-report.pdf?la=en>

Public Health Ontario. 2013. *Guide for Public Health Units: Considerations for Adult Mosquito Control*. Toronto, ON: Queen's Printer for Ontario.

Reisen, W.K., Y. Fang, and V.M. Martinez. 2006. Effects of temperature on the transmission of West Nile virus by *Culex tarsalis* (Diptera: Culicidae). *Journal of Medical Entomology* 43: 309-317.

Russell, C. 2008. *Analysis of the Feeding Behaviour of the Mosquito Culex pipiens L. (Diptera: Culicidae) in Relation to West Nile Virus*. Ph.D. Thesis, Brock University, St. Catharines, Ontario.

Turell, M.J., M.L. O'Guinn, D.J. Dohm, and J.W. Jones. 2001. Vector competence of North American mosquitoes (Diptera: Culicidae) for West Nile virus. *J. Med. Ent.* 38: 130-134.

Wood, D.M., P.T. Dang, and R.A. Ellis. 1979. *The Insects and Arachnids of Canada: Part 6. The Mosquitoes of Canada (Diptera: Culicidae)*. Biosystematics Research Institute, Ottawa.

